Abstract Submitted to the International Conference on Strongly Correlated Electron Systems University of Michigan, Ann Arbor August 6-10, 2001

VORTEX PHASE DIAGRAMS IN R_3 **Rh**₄**Sn**₁₃ (R = **Yb**, **Ca**)

C.V. Tomy¹, S. Sarkar², S. Ramakrishnan², A. K. Grover², Geetha Balakrishnan³, D. McK. Paul³

- ¹ Department of Physics, Indian Institute of Technology, Bombay 400 076, India
- ² Tata Institute of Fundamental Research, Mumbai-400 005, India
- ³ Department of Physics, University of Warwick, Coventry CV4 7AL, UK

The weakly pinned, isotropic low T_c superconductors $Yb_3Rh_4Sn_{13}$ and $Ca_3Rh_4Sn_{13}$ have emerged as important systems to witness a variety of issues related to phase transitions in vortex matter [1,2]. The (soft) vortex medium is a convenient prototype for studying the interplay as well as competition between interaction and disorder effects in condensed matter. In weakly pinned systems, the critical current density J_c imbibes information on the spatial and temporal correlation in the flux line lattice (FLL). The anomalous variations in $J_c(H,T)$ thus provide information on the transformations of the vortex matter. We present a status report on the studies related to the two well-known anomalous variations in $J_c(H,T)$ known as the Peak Effect (PE) and the Second Magnetization Peak (SMP) in $R_3Rh_4Sn_{13}$ (R=Yb, Ca) crystals. The novel features include the pinning induced step-wise amorphization of the FLL and the observation of the SMP and the PE in an isothermal scan. The close resemblances between the observed behavior in these compounds and those in anisotropic high T_c YBa $_2$ Cu $_3$ O $_7$ and the low T_c 2H-NbSe $_2$ shall be highlighted.

1. C.V. Tomy *et al* Physica C **280**, 1 (1997); Phys. Rev. B **56**, 8346 (1997). 2. S. Sarkar *et al* Phys. Rev. B **61**, 12394 (2000); Physica C **341-348**, 1055 (2000);**341-348**, 1085 (2000); Physica C (in press).